

2011 Paper 3

2011.3.1	Question 1		34
2011.3.2	Question 2		35
2011.3.3	Question 3		36
2011.3.4	Question 4		38
2011.3.5	Question 5		40
2011.3.6	Question 6		42
2011.3.7	Question 7		44
2011.3.8	Question 8		46
2011.3.12	Question 12		48
2011.3.13	Question 13		50

2011.3 Question 1

1. By rearrangement, we have

$$\frac{du}{u} = \left(1 + \frac{1}{x+1}\right) dx,$$

and hence by integration,

$$\ln|u| = x + \ln|x+1| + C.$$

This gives

$$u = C(x+1)e^x$$

as the general solution.

2. Since $y = ze^{-x}$, we must have

$$\frac{dy}{dx} = \frac{dz}{dx}e^{-x} - ze^{-x} = \left(\frac{dz}{dx} - z\right)e^{-x},$$

and

$$\frac{d^2y}{dx^2} = \frac{d^2z}{dx^2}e^{-x} - 2\frac{dz}{dx}e^{-x} + ze^{-x} = \left(\frac{d^2z}{dx^2} - 2\frac{dz}{dx} + z\right)e^{-x}.$$

Hence, the original differential equation can be simplified:

$$\begin{aligned} (x+1)\frac{d^2y}{dx^2} + x\frac{dy}{dx} - y &= 0 \\ (x+1)\left(\frac{d^2z}{dx^2} - 2\frac{dz}{dx} + z\right)e^{-x} + x\left(\frac{dz}{dx} - z\right)e^{-x} - ze^{-x} &= 0 \\ (x+1)\left(\frac{d^2z}{dx^2} - 2\frac{dz}{dx} + z\right) + x\left(\frac{dz}{dx} - z\right) - z &= 0 \\ (x+1)\frac{d^2z}{dx^2} - (x+2)\frac{dz}{dx} &= 0, \end{aligned}$$

which is a first-order differential equation for $\frac{dz}{dx}$.

Hence, from part (i), we have the general solution to this differential equation is

$$\frac{dz}{dx} = C(x+1)e^x,$$

and hence by integration

$$z = C \int (x+1)e^x dx = C \left[\int x de^x + \int e^x dx \right] = C[xe^x - e^x + e^x] + D.$$

Therefore, $y = ze^{-x} = De^{-x} + Cx$. Let $A = C$ and $B = D$ and this is exactly what is desired.

3. The complementary function is the differential equation solved in the previous part. For the complementary function, consider $y = ax^2 + b$, and hence $\frac{dy}{dx} = 2ax$ and $\frac{d^2y}{dx^2} = 2a$. Hence,

$$2a(x+1) + x \cdot 2ax - ax^2 - c = ax^2 + 2ax + (2a - c) = x^2 + 2x + 1.$$

Hence, $a = 1$ and $c = 1$ giving $y = x^2 + 1$ is a particular integral.

Therefore, the general solution to the differential equation is

$$y = Ax + Be^{-x} + x^2 + 1.$$

2011.3 Question 2

By definition,

$$f(x) = \sum_{k=0}^n a_k x^k$$

where $a_n = 1$.

Hence,

$$\begin{aligned} q^{n-1} f\left(\frac{p}{q}\right) &= q^{n-1} \sum_{k=0}^n a_k \left(\frac{p}{q}\right)^k \\ &= q^{n-1} \sum_{k=0}^n a_k p^k q^{-k} \\ &= \sum_{k=0}^n a_k p^k q^{n-k-1}. \end{aligned}$$

For the terms with $k = 0, 1, 2, \dots, n-1$, we have $n-k-1 \geq 0$ and hence the terms $a_k p^k q^{n-k-1}$ is an integer, and hence the sum from $k = 0$ to $k = n-1$ is an integer as well.

If $\frac{p}{q}$ is a rational root of f , $f\left(\frac{p}{q}\right) = 0$, and since all the rest of the terms are integers, the term where $k = n$ must be an integer as well. When $k = n$,

$$a_n p^n q^{n-n-1} = a_n p^n q^{-1} = \frac{p^n}{q}$$

must be an integer. But since p and q are co-prime, this can be an integer if and only if $q = 1$.

Therefore, $\frac{p}{q} = p$ is an integer as well, and any rational root to $f(x) = 0$ must be an integer.

1. Consider the polynomial $f(x) = x^n - 2$. The n th root of 2 must satisfy $1 < \sqrt[n]{2} < 2$, for $n \geq 2$. This is because $1^n = 1 < 2$ and $2^n = 2 \cdot 2^{n-1} > 21 = 2$.

The n th root of 2 is a root to f . If it is rational, then it must be integer. But $1 < \sqrt[n]{2} < 2$ and so the n th root of 2 cannot be an integer. Therefore, it must be irrational.

2. Consider the polynomial $f(x) = x^3 - x + 1$. If the roots to this polynomial are rational, then they must be integer.

Under modulo 2, $x^3 \equiv x$ since $1^3 \equiv 1$ and $0^3 \equiv 0$. Hence, $f(x) \equiv x^3 - x + 1 \equiv 0 + 1 \equiv 1$ modulo 2. This means there is no integer root to $f(x) = 0$ since the right-hand side is congruent to 0 modulo 2, and hence there are no rational roots.

3. Consider the polynomial $f(x) = x^n - 5x + 7$. If the roots to this polynomial are rational, then they must be integer.

For $n \geq 2$, under modulo 2, $x^n \equiv 5x$ since $1^n \equiv 1 \equiv 5 \equiv 5 \cdot 1$ and $0^n \equiv 0 \equiv 5 \cdot 0$. Hence, $f(x) \equiv x^n - 5x + 7 \equiv 0 + 7 \equiv 1$ modulo 2. This means there is no integer root to $f(x) = 0$ since the right-hand side is congruent to 0 modulo 2, and hence there are no rational roots.

2011.3 Question 3

We have

$$\begin{aligned} a(x - \alpha)^3 + b(x - \beta)^3 &= ax^3 - 3a\alpha x^2 + 3a\alpha^2 x - a\alpha^3 + bx^3 - 3b\beta x^2 + 3b\beta^2 x - b\beta^3 \\ &= (a + b)x^3 - 3(a\alpha + b\beta)x^2 + 3(a\alpha^2 + b\beta^2)x - (a\alpha^3 + b\beta^3). \end{aligned}$$

By comparing coefficients, we have

$$\begin{cases} a + b = 1, \\ -3(a\alpha + b\beta) = 0 \implies a\alpha + b\beta = 0, \\ 3(a\alpha^2 + b\beta^2) = -3p \implies a\alpha^2 + b\beta^2 = -p, \\ -(a\alpha^3 + b\beta^3) = q \implies a\alpha^3 + b\beta^3 = -q. \end{cases}$$

The first pair of equation solve to

$$(a, b) = \left(-\frac{\beta}{\alpha - \beta}, \frac{\alpha}{\alpha - \beta} \right).$$

Putting this into the third equation, we can see

$$\begin{aligned} \text{LHS} &= \frac{\beta}{\beta - \alpha} \cdot \alpha^2 - \frac{\alpha}{\beta - \alpha} \cdot \beta^2 \\ &= \frac{\alpha\beta(\alpha - \beta)}{\beta - \alpha} \\ &= -\alpha\beta \\ &= -\frac{p^2}{p} \\ &= -p \\ &= \text{RHS}, \end{aligned}$$

using Vieta's Theorem for $\alpha\beta$, and for the final one,

$$\begin{aligned} \text{LHS} &= \frac{\beta}{\beta - \alpha} \cdot \alpha^3 - \frac{\alpha}{\beta - \alpha} \cdot \beta^3 \\ &= \frac{\alpha\beta(\alpha^2 - \beta^2)}{\beta - \alpha} \\ &= -\frac{\alpha\beta(\alpha + \beta)(\beta - \alpha)}{\beta - \alpha} \\ &= -\alpha\beta(\alpha + \beta) \\ &= -\frac{p^2}{p} \cdot \left(-\frac{q}{p} \right) \\ &= -p \cdot \frac{q}{p} \\ &= -q \\ &= \text{RHS}, \end{aligned}$$

using Vieta's Theorem for $\alpha\beta$ and $\alpha + \beta$. Hence, this means for α, β being solutions to $pt^2 - qt + p^2 = 0$ and

$$(a, b) = \left(-\frac{\beta}{\alpha - \beta}, \frac{\alpha}{\alpha - \beta} \right),$$

we have

$$x^3 - 3px + q = a(x - \alpha)^3 + b(x - \beta)^3.$$

In this case here, we have $p = 8$ and $q = 48$. Hence, the quadratic equation is

$$8t^2 - 48t + 8^2 = 8(t^2 - 6t + 8) = 8(t - 2)(t - 4) = 0,$$

which solves to $(\alpha, \beta) = (2, 4)$ or $(\alpha, \beta) = (4, 2)$. Without loss of generality, let $(\alpha, \beta) = (2, 4)$, and hence

$$(a, b) = \left(-\frac{\beta}{\alpha - \beta}, \frac{\alpha}{\alpha - \beta} \right) = \left(-\frac{4}{2 - 4}, \frac{2}{2 - 4} \right) = (2, -1),$$

Hence, the original cubic equation

$$x^3 - 24x + 48 = 0$$

can be simplified to

$$2(x - 2)^3 - (x - 4)^3 = 0.$$

Hence,

$$2(x - 2)^3 = (x - 4)^3,$$

and we have

$$2^{\frac{1}{3}}(x - 2) = \omega^n(x - 4),$$

for $n = 0, 1, 2$ and $\omega = \exp\left(\frac{2\pi i}{3}\right)$.

Rearranging gives us

$$x = \frac{2\left(2\omega^n - 2^{\frac{1}{3}}\right)}{\omega^n - 2^{\frac{1}{3}}}$$

When $n = 0$, $\omega^n = 1$, and hence

$$x = \frac{2\left(2 - 2^{\frac{1}{3}}\right)}{1 - 2^{\frac{1}{3}}}.$$

The other two solutions

$$x = \frac{2\left(2\omega - 2^{\frac{1}{3}}\right)}{\omega - 2^{\frac{1}{3}}}, x = \frac{2\left(2\omega^2 - 2^{\frac{1}{3}}\right)}{\omega^2 - 2^{\frac{1}{3}}}.$$

This equation reduces to

$$x^3 - 3r^2x + 2r^3 = 0.$$

This can be factorised to


$$(x - r)(x^2 + rx - 2r^2) = (x - r)^2(x + 2r)$$

and the solutions are

$$x_{1,2} = r, x_3 = -2r.$$

2011.3 Question 4

1. The following two diagrams shows the cases $a < b$ and $a > b$ respectively.



In both cases, the shaded area is greater than the area of the rectangle formed by $(0,0)$, $(a,0)$, (a,b) and $(0,b)$, leading to the inequality. The equal sign holds when $b = f(a)$.

2. Since $f(x) = x^{p-1}$, we must have $x = f^{-1}(x)^{p-1}$, and hence $f^{-1}(x) = x^{\frac{1}{p-1}}$. Hence,

$$\int_0^a f(x) dx = \frac{1}{p} [x^p]_0^a = \frac{a^p}{p}.$$

Since $\frac{1}{p} + \frac{1}{q} = 1$, we must have $\frac{1}{q} = 1 - \frac{1}{p} = \frac{p-1}{p}$, and

$$q = \frac{p}{p-1},$$

and hence

$$f^{-1}(x) = x^{q-1},$$

which gives

$$\int_0^b f^{-1}(x) dx = \frac{b^q}{q}.$$

Since f is a polynomial, it must be continuous. $f(0) = 0^{p-1} = 0$, and

$$f'(x) = (p-1)x^{p-2}$$

is always non-negative for $x \geq 0$, we must have by the original inequality

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}$$

as desired.

3. Consider the function $f(x) = \sin x$. First, f is continuous, and

$$f'(x) = \cos x$$

is always positive for $0 \leq x \leq \frac{1}{2}\pi$. We notice

$$\int_0^a f(x) dx = [-\cos x]_0^a = 1 - \cos a,$$

and $f^{-1}(x) = \arcsin x$, and hence for $0 \leq b \leq 1$,

$$\begin{aligned} \int_0^b f^{-1}(x) \, dx &= \int_0^b \arcsin(x) \, dx \\ &= [x \arcsin x]_0^b - \int_0^b x \cdot \frac{1}{\sqrt{1-x^2}} \, dx \\ &= \left[x \arcsin x + \sqrt{1-x^2} \right]_0^b \\ &= b \arcsin b + \sqrt{1-b^2} - 1. \end{aligned}$$

Hence, using the given inequality,

$$ab \leq b \arcsin b + \sqrt{1-b^2} - 1 + 1 - \cos a = b \arcsin b + \sqrt{1-b^2} - \cos a,$$

as desired.

Let $a = 0$ and $b = t^{-1}$. Since $t \geq 1$, we have $0 < b \leq 1$, and hence

$$0 \leq t^{-1} \arcsin t^{-1} + \sqrt{1-t^{-2}} - \cos 0.$$

Multiplying both sides by t , and noticing $\cos 0 = 1$, we have

$$0 \leq \arcsin t^{-1} + \sqrt{t^2 - 1} - t,$$

and hence

$$\arcsin t^{-1} \geq t - \sqrt{t^2 - 1},$$

as desired.

2011.3 Question 5

Since we have

$$\tan \theta = \frac{y}{x} \implies \theta = \arctan \frac{y}{x} + k\pi$$

for some $k \in \mathbb{Z}$, differentiating with respect to t gives us

$$\frac{d\theta}{dt} = \frac{1}{1 + \left(\frac{y}{x}\right)^2} \cdot \frac{d\frac{y}{x}}{dt} = \frac{x^2}{x^2 + y^2} \cdot \frac{x \frac{dy}{dt} - y \frac{dx}{dt}}{x^2} = \frac{x \frac{dy}{dt} - y \frac{dx}{dt}}{r^2}.$$

Hence,

$$\frac{1}{2} \int r^2 d\theta = \frac{1}{2} \int r^2 \cdot \frac{x \frac{dy}{dt} - y \frac{dx}{dt}}{r^2} \cdot dt = \frac{1}{2} \int \left(x \frac{dy}{dt} - y \frac{dx}{dt} \right) dt,$$

as desired.

The coordinates of A and B are

$$A(x - a \cos t, y - a \sin t), B(x + b \cos t, y + b \sin t).$$

Hence, we have

$$\begin{aligned} [A] &= \frac{1}{2} \int_0^{2\pi} \left(x_A \frac{dy_A}{dt} - y_A \frac{dx_A}{dt} \right) dt \\ &= \frac{1}{2} \int_0^{2\pi} \left[(x - a \cos t) \left(\frac{dy}{dt} - a \cos t \right) - (y - a \sin t) \left(\frac{dx}{dt} + a \sin t \right) \right] dt \\ &= \frac{1}{2} \int_0^{2\pi} \left(x \frac{dy}{dt} - y \frac{dx}{dt} \right) dt - \frac{a}{2} \int_0^{2\pi} \left[\cos t \left(\frac{dy}{dt} + x \right) + \sin t \left(y - \frac{dx}{dt} \right) \right] dt \\ &\quad + \frac{a^2}{2} \int_0^{2\pi} (\cos^2 t + \sin^2 t) dt \\ &= [P] - af + \frac{a^2}{2} \int_0^{2\pi} dt \\ &= [P] - af + 2\pi \cdot \frac{a^2}{2} \\ &= [P] + \pi a^2 - af, \end{aligned}$$

as desired.

Similarly,

$$\begin{aligned} [B] &= \frac{1}{2} \int_0^{2\pi} \left(x_B \frac{dy_B}{dt} - y_B \frac{dx_B}{dt} \right) dt \\ &= \frac{1}{2} \int_0^{2\pi} \left[(x + b \cos t) \left(\frac{dy}{dt} + b \cos t \right) - (y + b \sin t) \left(\frac{dx}{dt} - b \sin t \right) \right] dt \\ &= \frac{1}{2} \int_0^{2\pi} \left(x \frac{dy}{dt} - y \frac{dx}{dt} \right) dt + \frac{b}{2} \int_0^{2\pi} \left[\cos t \left(\frac{dy}{dt} + x \right) + \sin t \left(y - \frac{dx}{dt} \right) \right] dt \\ &\quad + \frac{b^2}{2} \int_0^{2\pi} (\cos^2 t + \sin^2 t) dt \\ &= [P] + bf + \frac{b^2}{2} \int_0^{2\pi} dt \\ &= [P] + bf + 2\pi \cdot \frac{b^2}{2} \\ &= [P] + \pi b^2 + bf. \end{aligned}$$

Since over $t \in [0, 2\pi]$, A and B both trace over $\mathcal{D}$, we must have

$$[A] = [B],$$

and hence

$$\pi a^2 - af = \pi b^2 + bf,$$

which means

$$\pi(a+b)(a-b) = (a+b)f,$$

and hence

$$f = (a-b)\pi,$$

and therefore

$$[A] = [B] = [P] + ab\pi.$$

The area between the curves $\mathcal{C}$ and $\mathcal{D}$ is represented as $[A] - [P]$ or $[B] - [P]$, and hence this area is πab , as desired.

2011.3 Question 6

We show that T is equal to each of U, V, X , and by transitivity, this shows that all four are equal.

- To show $T = U$, consider the substitution $u = 2 \operatorname{artanh} t$, and hence $t = \tanh \frac{u}{2}$.

When $t = \frac{1}{2}$, $u = 2 \operatorname{artanh} \frac{1}{2} = 2 \cdot \frac{1}{2} \cdot \ln \left(\frac{1+\frac{1}{2}}{1-\frac{1}{2}} \right) = \ln 3$, and when $t = \frac{1}{3}$, $u = 2 \operatorname{artanh} \frac{1}{3} = 2 \cdot \frac{1}{2} \cdot \ln \left(\frac{1+\frac{1}{3}}{1-\frac{1}{3}} \right) = \ln 2$.

We have $du = \frac{2}{1-t^2} dt$, and hence

$$\begin{aligned} T &= \int_{\frac{1}{3}}^{\frac{1}{2}} \frac{\operatorname{artanh} t}{t} dt \\ &= \int_{\ln 2}^{\ln 3} \frac{\frac{u}{2}}{\tanh \frac{u}{2}} \cdot \frac{1 - \tanh^2 \frac{u}{2}}{2} du \\ &= \int_{\ln 2}^{\ln 3} \frac{u}{2} \cdot \frac{1 - \tanh^2 \frac{u}{2}}{2 \tanh \frac{u}{2}} du \\ &= \int_{\ln 2}^{\ln 3} \frac{u}{2 \sinh u} du \\ &= U. \end{aligned}$$

- To show $T = V$, we use integration by parts.

$$\begin{aligned} T &= \int_{\frac{1}{3}}^{\frac{1}{2}} \frac{\operatorname{artanh} t}{t} dt \\ &= \int_{\frac{1}{3}}^{\frac{1}{2}} \operatorname{artanh} t \, d \ln t \\ &= [\operatorname{artanh} t \ln t]_{\frac{1}{3}}^{\frac{1}{2}} - \int_{\frac{1}{3}}^{\frac{1}{2}} \ln t \, d \operatorname{artanh} t \\ &= \left(\operatorname{artanh} \frac{1}{2} \ln \frac{1}{2} - \operatorname{artanh} \frac{1}{3} \ln \frac{1}{3} \right) - \int_{\frac{1}{3}}^{\frac{1}{2}} \frac{\ln t}{1-t^2} dt \\ &= \left(\frac{1}{2} \cdot \ln \left(\frac{1+\frac{1}{2}}{1-\frac{1}{2}} \right) \cdot (-\ln 2) - \frac{1}{2} \cdot \ln \left(\frac{1+\frac{1}{3}}{1-\frac{1}{3}} \right) \cdot (-\ln 3) \right) + V \\ &= \left(-\frac{1}{2} \cdot \ln 3 \cdot \ln 2 + \frac{1}{2} \cdot \ln 2 \cdot \ln 3 \right) + V \\ &= V. \end{aligned}$$

- To show $T = X$, consider the substitution $x = -\frac{1}{2} \ln t$, and hence $t = e^{-2x}$.

When $t = \frac{1}{2}$, $x = -\frac{1}{2} \ln \frac{1}{2} = \frac{1}{2} \ln 2$, and when $t = \frac{1}{3}$, $x = -\frac{1}{2} \ln \frac{1}{3} = \frac{1}{2} \ln 3$.

We have $dx = -\frac{dt}{2t}$, and hence

$$\begin{aligned} T &= \int_{\frac{1}{3}}^{\frac{1}{2}} \frac{\operatorname{artanh} t}{t} dt \\ &= \int_{\frac{1}{2} \ln 3}^{\frac{1}{2} \ln 2} \frac{\operatorname{artanh} e^{-2x}}{t} \cdot (-2t) dx \\ &= \int_{\frac{1}{2} \ln 2}^{\frac{1}{2} \ln 3} 2 \operatorname{artanh} e^{-2x} dx \\ &= \int_{\frac{1}{2} \ln 2}^{\frac{1}{2} \ln 3} \ln \left(\frac{1 + e^{-2x}}{1 - e^{-2x}} \right) dx \\ &= \int_{\frac{1}{2} \ln 2}^{\frac{1}{2} \ln 3} \ln \left(\frac{e^x + e^{-x}}{e^x - e^{-x}} \right) dx \\ &= \int_{\frac{1}{2} \ln 2}^{\frac{1}{2} \ln 3} \ln \coth x dx \\ &= X. \end{aligned}$$

2011.3 Question 7

1. The base case is when $n = 2$, and we have

$$T_2 = (\sqrt{a+1} + \sqrt{a})^2 = (2a+1) + 2 \cdot \sqrt{a(a+1)}.$$

We therefore have $A_2 = 2a+1$ and $B_2 = 2$, and we verify that

$$a(a+1)B_2^2 + 1 = a(a+1) \cdot 2^2 + 1 = 4a^2 + 4a + 1 = (2a+1)^2 = A_2^2,$$

as desired, and the statement holds for the base case when $n = 2$.

Now, assume that this statement is for some even $n = k$, i.e.

$$T_k = A_k + B_k \sqrt{a(a+1)}$$

where A_k and B_k are both integers, and $A_k^2 = a(a+1)B_k^2 + 1$.

Notice that

$$\begin{aligned} T_{k+2} &= T_k \cdot (\sqrt{a+1} + \sqrt{a})^2 \\ &= (A_k + B_k \sqrt{a(a+1)}) \cdot (2a+1 + 2\sqrt{a(a+1)}) \\ &= A_k \cdot (2a+1) + B_k \cdot 2 \cdot a(a+1) + 2A_k \sqrt{a(a+1)} + (2a+1)B_k \sqrt{a(a+1)} \\ &= [(2a+1)A_k + 2a(a+1)B_k] + [2A_k + (2a+1)B_k] \sqrt{a(a+1)}. \end{aligned}$$

Now let $A_{k+2} = (2a+1)A_k + 2a(a+1)B_k$, and $B_{k+2} = 2A_k + (2a+1)B_k$. Since a is a positive integer, and A_k and B_k are both integers, we must have A_{k+2} and B_{k+2} are both integers. Furthermore,

$$\begin{aligned} &A_{k+2}^2 - [a(a+1)B_{k+2}^2 + 1] \\ &= [(2a+1)A_k + 2a(a+1)B_k]^2 - [a(a+1)(2A_k + (2a+1)B_k)^2 + 1] \\ &= [(2a+1)^2 - 4a(a+1)]A_k^2 \\ &\quad + [2 \cdot (2a+1) \cdot 2a(a+1) - 2 \cdot a(a+1) \cdot 2 \cdot (2a+1)]A_k B_k \\ &\quad + [(2a(a+1))^2 - a(a+1)(2a+1)^2]B_k^2 - 1 \\ &= A_k^2 - a(a+1)B_k^2 - 1 \\ &= 1 - 1 \\ &= 0, \end{aligned}$$

and hence

$$A_{k+2}^2 = a(a+1)B_{k+2}^2 + 1.$$

So the original statement holds for $n = k+2$.

By the principle of mathematical induction, the original statement must hold for all even integers n .

2. If n is odd, then we have

$$\begin{aligned} T_n &= (\sqrt{a+1} + \sqrt{a})T_{n-1} \\ &= (\sqrt{a+1} + \sqrt{a})(A_{n-1} + B_{n-1}\sqrt{a(a+1)}) \\ &= A_{n-1}\sqrt{a+1} + A_{n-1}\sqrt{a} + B_{n-1}(a+1)\sqrt{a} + B_{n-1}a\sqrt{a+1} \\ &= (A_{n-1} + aB_{n-1})\sqrt{a+1} + (A_{n-1} + (a+1)B_{n-1})\sqrt{a}. \end{aligned}$$

Now, consider $C_n = A_{n-1} + aB_{n-1}$, and $D_n = A_{n-1} + (a+1)B_{n-1}$. Since a is a positive integer,

and A_{n-1} and B_{n-1} are integers, we must have C_n and D_n are integers as well. Furthermore,

$$\begin{aligned}
 & (a+1)C_n^2 - (aD_n^2 + 1) \\
 &= (a+1)(A_{n-1} + aB_{n-1})^2 - \left[a(A_{n-1} + (a+1)B_{n-1})^2 + 1 \right] \\
 &= [(a+1) - a]A_{n-1}^2 + [(a+1) \cdot 2 \cdot a - a \cdot 2 \cdot (a+1)]A_{n-1}B_{n-1} \\
 &\quad + [(a+1)a^2 - a(a+1)^2]B_{n-1}^2 - 1 \\
 &= A_{n-1}^2 - a(a+1)B_{n-1}^2 - 1 \\
 &= 1 - 1 \\
 &= 0,
 \end{aligned}$$

and hence

$$(a+1)C_n^2 = aD_n^2 + 1.$$

This shows precisely the original statement.

3. For even n ,

$$T_n = A_n + B_n\sqrt{a(a+1)} = \sqrt{A_n^2} + \sqrt{B_n^2 \cdot a(a+1)} = \sqrt{A_n^2} + \sqrt{A_n^2 - 1},$$

and for odd n ,

$$T_n = C_n\sqrt{a+1} + D_n\sqrt{a} = \sqrt{C_n^2(a+1)} + \sqrt{D_n^2 a} = \sqrt{aD_n^2 + 1} + \sqrt{aD_n^2},$$

as desired.

2011.3 Question 8

Since $w = u + iv$, $z = x + iy$, we have

$$\begin{aligned}
 u + iv &= w \\
 &= \frac{1 + iz}{i + z} \\
 &= \frac{1 + i(x + iy)}{i + (x + iy)} \\
 &= \frac{(1 - y) + xi}{x + (y + 1)i} \\
 &= \frac{(1 - y) + xi}{x + (y + 1)i} \cdot \frac{x - (y + 1)i}{x - (y + 1)i} \\
 &= \frac{[(1 - y) + xi][x - (y + 1)i]}{x^2 + (y + 1)^2} \\
 &= \frac{(1 - y)x + x(y + 1)}{x^2 + (y + 1)^2} + \frac{x^2 - (1 - y) \cdot (y + 1)}{x^2 + (y + 1)^2} \cdot i \\
 &= \frac{2x}{x^2 + (y + 1)^2} + \frac{x^2 + y^2 - 1}{x^2 + (y + 1)^2} \cdot i,
 \end{aligned}$$

and hence

$$(u, v) = \left(\frac{2x}{x^2 + (y + 1)^2}, \frac{x^2 + y^2 - 1}{x^2 + (y + 1)^2} \right).$$

1. When $y = 0$, we have

$$(u, v) = \left(\frac{2x}{x^2 + 1}, \frac{x^2 - 1}{x^2 + 1} \right).$$

Let $x = \tan\left(\frac{\theta}{2}\right)$. The tangent half-angle substitution also gives that $u = \sin \theta$ and $v = -\cos \theta$, and hence $u^2 + v^2 = 1$.

For the range of θ , we have $-\frac{\pi}{2} < \frac{\theta}{2} < \frac{\pi}{2}$, which means $-\pi < \theta < \pi$.

This represents the unit circle without the point $(\sin \pi, -\cos \pi) = (0, 1)$ corresponding to $\theta = \pi(+2k\pi)$ for some integer k .

2. When $-1 < x < 1$, we have $-\frac{\pi}{4} < \frac{\theta}{2} < \frac{\pi}{4}$, which means $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$. This is the unit circle with only the part below the u axis (exclusive).

3. When $x = 0$, we have

$$(u, v) = \left(0, \frac{y^2 - 1}{(y + 1)^2} \right).$$

Notice that

$$v = \frac{y^2 - 1}{(y + 1)^2} = \frac{(y + 1)(y - 1)}{(y + 1)^2} = \frac{y - 1}{y + 1} = 1 - \frac{2}{y + 1},$$

and hence $-1 < v < 1$.

This means the locus of w is the line segment $u = 0, -1 < v < 1$.

4. When $y = 1$, we have

$$(u, v) = \left(\frac{2x}{x^2 + 4}, \frac{x^2}{x^2 + 4} \right).$$

First, let $x = 2t$, and we have

$$(u, v) = \left(\frac{4t}{4t^2 + 4}, \frac{4t^2}{4t^2 + 4} \right) = \left(\frac{t}{t^2 + 1}, \frac{t^2}{t^2 + 1} \right).$$

Let $t = \tan\left(\frac{\theta}{2}\right)$, and we have $-\pi < \theta < \pi$. Notice that

$$u = \frac{1}{2} \cdot \frac{2t}{t^2 + 1} = \frac{1}{2} \sin \theta,$$

and

$$v - \frac{1}{2} = \frac{1}{2} \cdot \frac{t^2 - 1}{t^2 + 1} = -\frac{1}{2} \cos \theta.$$

This means the loci is a subset of the circle centred at $(0, \frac{1}{2})$ with radius $\frac{1}{2}$, with the point

$$(u, v) = \left(\frac{1}{2} \sin \pi, \frac{1}{2} - \frac{1}{2} \cos \pi \right) = (0, 1)$$

missing, which corresponds to $\theta = \pi(+2k\pi)$ for some integer k .

2011.3 Question 12

By differentiation, we have

$$[G(H(t))]' = G'(H(t)) \cdot H'(t).$$

Hence, we have

$$\begin{aligned} E(Y) &= [G(H(t))]'|_{t=1} \\ &= G'(H(1)) \cdot H'(1) \\ &= G'(1) \cdot H'(1) \\ &= E(N) \cdot E(X_i). \end{aligned}$$

By differentiating twice, we have

$$[G(H(t))]' = G''(H(t)) \cdot H'(t) \cdot H'(t) + G'(H(t)) \cdot H''(t).$$

Hence, we have

$$\begin{aligned} \text{Var}(Y) &= E(Y(Y-1)) + E(Y) - E(Y)^2 \\ &= [G(H(t))]'|_{t=1} + E(Y) - E(Y)^2 \\ &= G''(H(1)) \cdot H'(1) \cdot H'(1) + G'(H(1)) \cdot H''(1) + E(Y) - E(Y)^2 \\ &= G''(1) \cdot H'(1)^2 + G'(1) \cdot H''(1) + E(Y) - E(Y)^2 \\ &= E(N(N-1)) \cdot E(X_i)^2 + E(N) \cdot E(X_i(X_i-1)) + E(Y) - E(Y)^2 \\ &= [\text{Var}(N) + E(N)^2 - E(N)] \cdot E(X_i)^2 + E(N) \cdot [\text{Var}(X_i) + E(X_i^2) - E(X_i)] \\ &\quad + E(N) \cdot E(X_i) - E(N)^2 \cdot E(X_i)^2 \\ &= \text{Var}(N) E(X_i)^2 + E(N) \text{Var}(X_i). \end{aligned}$$

As defined, we have $N \sim \text{Geo}(\frac{1}{2})$, and hence

$$G(t) = \frac{\frac{1}{2} \cdot t}{1 - (1 - \frac{1}{2})t} = \frac{t}{2-t},$$

and

$$E(N) = 1/\frac{1}{2} = 2, \text{Var}(N) = \frac{1 - \frac{1}{2}}{(\frac{1}{2})^2} = 2.$$

We have $X_i \sim B(1, \frac{1}{2})$, and hence

$$H(t) = \frac{1}{2} \cdot t^0 + \frac{1}{2} \cdot t^1 = \frac{1}{2} \cdot (1+t),$$

and

$$E(X_i) = 1 \cdot \frac{1}{2} = \frac{1}{2}, \text{Var}(X_i) = 1 \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4}.$$

Hence, for $Y = \sum_{i=1}^N X_i$, we have

$$\text{p.g.f.}_Y(t) = G(H(t)) = \frac{\frac{1}{2}(1+t)}{2 - \frac{1}{2}(1+t)} = \frac{1+t}{3-t},$$

and by the formula for expectation and variance, we have

$$E(Y) = E(N) E(X_i) = 2 \cdot \frac{1}{2} = 1,$$

and

$$\text{Var}(Y) = \text{Var}(N) \cdot E(X_i)^2 + E(N) \cdot \text{Var}(X_i) = 2 \cdot \left(\frac{1}{2}\right)^2 + 2 \cdot \frac{1}{4} = 1.$$

By expressing the probability generating function of Y as a power series, we notice that

$$\begin{aligned}\text{p.g.f.}_Y(t) &= \frac{1+t}{3-t} \\&= -1 + \frac{4}{3-t} \\&= -1 + \frac{4}{3} \cdot \frac{1}{1-\frac{t}{3}} \\&= -1 + \frac{4}{3} \sum_{r=0}^{\infty} \left(\frac{t}{3}\right)^r \\&= -1 + \frac{4}{3} + \frac{4}{3} \sum_{r=1}^{\infty} 3^{-r} \cdot t^r \\&= \frac{1}{3} + \frac{4}{3} \sum_{r=1}^{\infty} 3^{-r} \cdot t^r,\end{aligned}$$

and hence

$$P(Y = y) = \begin{cases} \frac{1}{3}, & y = 0, \\ \frac{4}{3^{y+1}}, & \text{otherwise.} \end{cases}$$

2011.3 Question 13

1. We first find the expression given by the question.

$$\begin{aligned}
 \frac{P(X = r + 1)}{P(X = r)} &= \frac{\left(\frac{b}{n}\right)^{r+1} \left(\frac{n-b}{n}\right)^{k-r-1} \binom{k}{r+1}}{\left(\frac{b}{n}\right)^r \left(\frac{n-b}{n}\right)^{k-r} \binom{k}{r}} \\
 &= \frac{b/n}{(n-b)/n} \cdot \frac{\frac{k!}{(r+1)!(k-r-1)!}}{\frac{k!}{r!(k-r)!}} \\
 &= \frac{b}{n-b} \cdot \frac{r!(k-r)!}{(r+1)!(k-r-1)!} \\
 &= \frac{b}{n-b} \cdot \frac{k-r}{r+1} \\
 &= \frac{b}{n-b} \cdot \left(\frac{k+1}{r+1} - 1\right),
 \end{aligned}$$

and we can see that this decreases as r increases.

If the most probable number of black balls in the sample is unique (let it be r_0), then we have

$$P(X = r_0 + 1) < P(X = r_0) \iff \frac{P(X = r_0 + 1)}{P(X = r_0)} < 1,$$

and

$$P(X = r_0 - 1) < P(X = r_0) \iff \frac{P(X = r_0)}{P(X = r_0 - 1)} > 1,$$

This means r_0 is the minimal solution to the inequality

$$\frac{P(X = r + 1)}{P(X = r)} < 1.$$

This could be simplified to

$$\begin{aligned}
 \frac{P(X = r + 1)}{P(X = r)} &< 1 \\
 \frac{b}{n-b} \left(\frac{k+1}{r+1} - 1\right) &< 1 \\
 \frac{k+1}{r+1} - 1 &< \frac{n-b}{b} \\
 \frac{k+1}{r+1} &< \frac{n}{b} \\
 r+1 &> \frac{b(k+1)}{n} \\
 r &> \frac{b(k+1)}{n} - 1,
 \end{aligned}$$

and hence

$$r_0 = \left\lfloor \frac{b(k+1)}{n} \right\rfloor.$$

It is not unique when there exists some r where

$$\frac{P(X = r_0 + 1)}{P(X = r_0)} = 1,$$

which means there exists an integer r such that

$$r = \frac{b(k+1)}{n} - 1.$$

This happens if and only if $n \mid b(k+1)$.

2. Let Y be the number of black balls in the sample. Similarly, we have

$$\begin{aligned}
 \frac{P(Y = r+1)}{P(Y = r)} &= \frac{\frac{\binom{b}{r+1} \cdot \binom{n-b}{k-r-1}}{\binom{n}{k}}}{\frac{\binom{b}{r} \cdot \binom{n-b}{k-r}}{\binom{n}{k}}} \\
 &= \frac{\frac{b!}{(r+1)!(b-r-1)!} \cdot \frac{(n-b)!}{(k-r-1)!(n+r-k-b+1)!}}{\frac{b!}{r!(b-r)!} \cdot \frac{(n-b)!}{(k-r)!(n+r-k-b)!}} \\
 &= \frac{r!(b-r)!(k-r)!(n+r-k-b)!}{(r+1)!(b-r-1)!(k-r-1)!(n+r-k-b+1)!} \\
 &= \frac{(b-r) \cdot (k-r)}{(r+1) \cdot (n+r-k-b+1)}.
 \end{aligned}$$

The most probable number of black balls is the smallest solution to

$$\begin{aligned}
 \frac{(b-r) \cdot (k-r)}{(r+1) \cdot (n+r-k-b+1)} &< 1 \\
 (b-r)(k-r) &< (r+1)(n+r-k-b+1) \\
 bk - rk - bk + r^2 &< nr + r^2 - rk - bk + r + n + r - k - b + 1 \\
 (n+2)r &> bk + k + b - 1 - n \\
 r &> \frac{bk + k + b - 1 - n}{n+2} \\
 &= \frac{(n+1)(k+1)}{n+2} - 1.
 \end{aligned}$$

This means the most probable number of black balls, given its uniqueness, is

$$\left\lfloor \frac{(b+1)(k+1)}{(n+2)} \right\rfloor.$$

It is not unique when

$$\frac{(n+1)(k+1)}{n+2} - 1$$

is an integer, if and only if

$$(n+2) \mid (n+1)(k+1).$$