

2017 Paper 3

2017.3.1	Question 1	31
2017.3.2	Question 2	33
2017.3.3	Question 3	34
2017.3.4	Question 4	35
2017.3.5	Question 5	37
2017.3.6	Question 6	39
2017.3.7	Question 7	42
2017.3.8	Question 8	45
2017.3.12	Question 12	47
2017.3.13	Question 13	49

2017.3 Question 1

1. We have

$$\begin{aligned}
 \text{RHS} &= \frac{r+1}{r} \left(\frac{1}{\binom{n+r-1}{r}} - \frac{1}{\binom{n+r}{r}} \right) \\
 &= \frac{r+1}{r} \left(\frac{r!(n-1)!}{(n+r-1)!} - \frac{r!n!}{(n+r)!} \right) \\
 &= \frac{r+1}{r} \left(\frac{r!(n-1)!(n+r)}{(n+r)!} - \frac{r!(n-1)!n}{(n+r)!} \right) \\
 &= \frac{r+1}{r} \cdot \frac{r!(n-1)!(n+r) - r!(n-1)!n}{(n+r)!} \\
 &= \frac{r+1}{r} \cdot \frac{r!(n-1)!r}{(n+r)!} \\
 &= \frac{(r+1)!(n-1)!}{(n+r)!} \\
 &= \binom{n+r}{r+1} \\
 &= \text{LHS}
 \end{aligned}$$

as desired.

Therefore,

$$\begin{aligned}
 \sum_{n=1}^{+\infty} \frac{1}{\binom{n+r}{r+1}} &= \sum_{n=1}^{+\infty} \frac{r+1}{r} \left(\frac{1}{\binom{n+r-1}{r}} - \frac{1}{\binom{n+r}{r}} \right) \\
 &= \frac{r+1}{r} \sum_{n=1}^{+\infty} \left(\frac{1}{\binom{n+r-1}{r}} - \frac{1}{\binom{n+r}{r}} \right) \\
 &= \frac{r+1}{r} \left[\sum_{n=0}^{+\infty} \frac{1}{\binom{n+r}{r}} - \sum_{n=1}^{+\infty} \frac{1}{\binom{n+r}{r}} \right] \\
 &= \frac{r+1}{r} \frac{1}{\binom{0+r}{r}} \\
 &= \frac{r+1}{r},
 \end{aligned}$$

assuming the sum converges.

When $r = 2$, we have

$$\sum_{n=1}^{+\infty} \frac{1}{\binom{n+2}{3}} = \frac{3}{2}.$$

When $n = 1$, $\frac{1}{\binom{1+2}{3}} = \frac{1}{1} = 1$.

Therefore,

$$\sum_{n=2}^{+\infty} \frac{1}{\binom{n+2}{3}} = \frac{1}{2}$$

as desired.

2. Notice that

$$\begin{aligned}
 \frac{3!}{n^3} < \frac{1}{\binom{n+1}{3}} &\iff \frac{3!}{n^3} < \frac{3!}{(n+1)n(n-1)} \\
 &\iff n^3 > (n+1)n(n-1) \\
 &\iff n^3 > n(n^2 - 1) \\
 &\iff n^3 > n^3 - n \\
 &\iff n > 0,
 \end{aligned}$$

which is true.

Also, notice that

$$\begin{aligned}
 \frac{20}{\binom{n+1}{3}} - \frac{1}{\binom{n+2}{5}} < \frac{5!}{n^3} &\iff \frac{5!}{(n+1)(n)(n-1)} - \frac{5!}{(n+2)(n+1)(n)(n-1)(n-2)} < \frac{5!}{n^3} \\
 &\iff \frac{(n+2)(n-2) - 1}{(n+2)(n+1)(n)(n-1)(n-2)} < \frac{1}{n^3} \\
 &\iff (n^2 - 5)n^3 < (n^2 - 4)(n^2 - 1)n \\
 &\iff n^5 - 5n^3 < n^5 - 5n^3 + 4n \\
 &\iff 4n > 0,
 \end{aligned}$$

which is true.

Therefore, we have that

$$\begin{aligned}
 \sum_{n=3}^{+\infty} \frac{3!}{n^3} &< \sum_{n=3}^{+\infty} \frac{1}{\binom{n+1}{3}} \\
 &= \sum_{n=2}^{+\infty} \frac{1}{\binom{n+2}{3}} \\
 &= \frac{1}{2},
 \end{aligned}$$

and therefore $\sum_{n=3}^{+\infty} \frac{1}{n^3} < \frac{1}{12}$, and $\sum_{n=1}^{+\infty} \frac{1}{n^3} < 1 + \frac{1}{8} + \frac{1}{12} = \frac{29}{24} = \frac{116}{96}$.

On the other hand, we have

$$\begin{aligned}
 \sum_{n=3}^{+\infty} \frac{5!}{n^3} &< \sum_{n=3}^{+\infty} \left[\frac{20}{\binom{n+1}{3}} - \frac{1}{\binom{n+2}{5}} \right] \\
 &= 20 \sum_{n=2}^{+\infty} \frac{1}{\binom{n+2}{3}} - \sum_{n=1}^{+\infty} \frac{1}{\binom{n+4}{5}} \\
 &= 20 \cdot \frac{1}{2} - \frac{5}{4} \\
 &= 10 - \frac{5}{4} \\
 &= \frac{35}{4},
 \end{aligned}$$

and therefore $\sum_{n=3}^{+\infty} \frac{1}{n^3} > \frac{7}{96}$, and $\sum_{n=1}^{+\infty} \frac{1}{n^3} > 1 + \frac{1}{8} + \frac{7}{96} = \frac{115}{96}$.

Hence,

$$\frac{115}{96} < \sum_{n=1}^{+\infty} \frac{1}{n^3} < \frac{116}{96}$$

as desired.

2017.3 Question 2

1. Let the complex number representing $R(P)$ be z' . Therefore,

$$\begin{aligned} z' - a &= \exp(i\theta)(z - a), \\ z' &= z \exp(i\theta) + a(1 - \exp(i\theta)), \end{aligned}$$

as desired.

2. Let the complex number representing $SR(P)$ be z'' . Therefore,

$$\begin{aligned} z'' - b &= \exp(i\varphi)(z' - b), \\ z'' &= z' \exp(i\varphi) + b(1 - \exp(i\varphi)), \\ z'' &= [z \exp(i\theta) + a(1 - \exp(i\theta))] \exp(i\varphi) + b(1 - \exp(i\varphi)), \\ z'' &= z \exp(i(\theta + \varphi)) + a(1 - \exp(i\theta)) \exp(i\varphi) + b(1 - \exp(i\varphi)). \end{aligned}$$

This will be an anti-clockwise rotation around c over an angle of $(\theta + \varphi)$, where

$$c[1 - \exp(i(\theta + \varphi))] = a \exp(i\varphi) - a \exp(i(\theta + \varphi)) + b - b \exp(i\varphi),$$

If $\theta + \varphi = 2n\pi$ for some integer $n \in \mathbb{Z}$, $1 - \exp(i(\theta + \varphi)) = 0$, therefore c cannot be determined.

Multiplying both sides by $\exp\left(-\frac{i(\theta + \varphi)}{2}\right)$, we have

$$\begin{aligned} &c \left[\exp\left(-\frac{i(\theta + \varphi)}{2}\right) - \exp\left(\frac{i(\theta + \varphi)}{2}\right) \right] \\ &= a \left[\exp\left(\frac{i(\varphi - \theta)}{2}\right) - \exp\left(\frac{i(\theta + \varphi)}{2}\right) \right] + b \left[\exp\left(-\frac{i(\theta + \varphi)}{2}\right) - \exp\left(\frac{i(\varphi - \theta)}{2}\right) \right], \end{aligned}$$

and hence

$$\begin{aligned} -2ci \sin\left(\frac{\theta + \varphi}{2}\right) &= -2ai \exp\left(\frac{i\varphi}{2}\right) \sin\left(\frac{\theta}{2}\right) - 2bi \exp\left(-\frac{i\theta}{2}\right) \sin\left(\frac{\varphi}{2}\right), \\ c \sin\left(\frac{\theta + \varphi}{2}\right) &= a \exp\left(\frac{i\varphi}{2}\right) \sin\left(\frac{\theta}{2}\right) + b \exp\left(-\frac{i\theta}{2}\right) \sin\left(\frac{\varphi}{2}\right). \end{aligned}$$

If $\theta + \varphi = 2\pi$, we will have $z'' = z + a \exp(i\varphi) - a + b(1 - \exp(i\varphi)) = z + (b - a)(1 - \exp(i\varphi))$, which is a translation by $(b - a)(1 - \exp(i\varphi))$.

3. If $RS = SR$, then we have

$$\begin{aligned} a(1 - \exp(i\theta)) \exp(i\varphi) + b(1 - \exp(i\varphi)) &= b(1 - \exp(i\varphi)) \exp(i\theta) + a(1 - \exp(i\theta)), \\ a(-1 + \exp(i\varphi) + \exp(i\theta) - \exp(i(\theta + \varphi))) &= b(-1 + \exp(i\varphi) + \exp(i\theta) - \exp(i(\theta + \varphi))), \\ (a - b)(1 - \exp(i\varphi))(1 - \exp(i\theta)) &= 0. \end{aligned}$$

Therefore, $a = b$, or $\varphi = 2n\pi$, or $\theta = 2n\pi$, for some integer $n \in \mathbb{Z}$.

2017.3 Question 3

By Vieta's Theorem, from the quartic equation in x , we have

$$\alpha\beta + \alpha\gamma + \alpha\delta + \beta\gamma + \beta\delta + \gamma\delta = q,$$

and from the cubic equation in y , we have

$$(\alpha\beta + \gamma\delta) + (\alpha\gamma + \beta\delta) + (\alpha\delta + \beta\gamma) = -A.$$

Therefore, $A = -q$.

1. Since $(p, q, r, s) = (0, 3, -6, 10)$, the cubic equation is reduced to

$$y^3 - 3y^2 - 10y + 84 = 0,$$

and therefore

$$(y - 2)(y - 7)(y + 6) = 0.$$

Therefore, $y_1 = 7, y_2 = 2, y_3 = -6$, and $\alpha\beta + \gamma\delta = 7$.

2. We have

$$\begin{aligned} (\alpha + \beta)(\gamma + \delta) &= \alpha\gamma + \alpha\delta + \beta\gamma + \beta\delta \\ &= (\alpha\beta + \alpha\gamma + \alpha\delta + \beta\gamma + \beta\delta + \gamma\delta) - (\alpha\beta + \gamma\delta) \\ &= q - 7 \\ &= 3 - 7 \\ &= -4. \end{aligned}$$

By Vieta's Theorem, we have $\alpha\beta\gamma\delta = s = 10$. Therefore, $\alpha\beta$ and $\gamma\delta$ must be roots to the equation

$$x^2 - 7x + 10 = 0.$$

The two roots are $x = 2$ and $x = 5$, and therefore $\alpha\beta = 5$.

3. We have from the other root that $\gamma\delta = 2$.

We notice that $(\alpha + \beta) + (\gamma + \delta) = -p = 0$. Therefore, from part 2, $(\alpha + \beta)$ and $(\gamma + \delta)$ are roots to the equation

$$x^2 - 4 = 0.$$

This gives us $\alpha + \beta = \pm 2$ and $\gamma + \delta = \mp 2$.

Using the value of r and Vieta's Theorem, we have

$$\alpha\beta\gamma + \alpha\beta\delta + \alpha\gamma\delta + \beta\gamma\delta = -r = 6.$$

Plugging in $\alpha\beta = 5$ and $\gamma\delta = 2$, we have

$$5(\gamma + \delta) + 2(\alpha + \beta) = 6.$$

Therefore, it must be the case that $\alpha + \beta = -2$ and $\gamma + \delta = 2$.

Hence, using the values of $\alpha\beta$ and $\gamma\delta$, α and β are solutions to the quadratic equation $x^2 + 2x + 5 = 0$, and γ and δ are solutions to the quadratic equation $x^2 - 2x + 2 = 0$.

Solving this gives us $\alpha, \beta = -1 \pm 2i$ and $\gamma, \delta = 1 \pm i$. The solutions to the original quartic equation is

$$x_{1,2} = -1 \pm 2i, x_{3,4} = 1 \pm i.$$

2017.3 Question 4

1. Notice that $a = e^{\ln a}$ and hence $a^x = e^{x \ln a}$, $a^{\frac{x}{\ln a}} = e^x$ we have

$$\begin{aligned} F(y) &= \exp\left(\frac{1}{y} \int_0^y \ln f(x) \, dx\right) \\ &= a^{\frac{1}{y \ln a} \cdot \int_0^y \ln f(x) \, dx} \\ &= a^{\frac{1}{y} \cdot \int_0^y \frac{\ln f(x)}{\ln a} \, dx} \\ &= a^{\frac{1}{y} \cdot \int_0^y \log_a f(x) \, dx} \end{aligned}$$

as desired.

2. We have

$$\begin{aligned} H(y) &= \exp\left(\frac{1}{y} \int_0^y \ln f(x)g(x) \, dx\right) \\ &= \exp\left[\frac{1}{y} \int_0^y (\ln f(x) + \ln g(x)) \, dx\right] \\ &= \exp\left[\frac{1}{y} \left(\int_0^y \ln f(x) \, dx + \int_0^y \ln g(x) \, dx\right)\right] \\ &= \exp\left(\frac{1}{y} \int_0^y \ln f(x) \, dx\right) \cdot \exp\left(\frac{1}{y} \int_0^y \ln g(x) \, dx\right) \\ &= F(y) \cdot G(y). \end{aligned}$$

3. Let $f(x) = b^x$.

$$\begin{aligned} F(y) &= \exp\left(\frac{1}{y} \int_0^y \ln f(x) \, dx\right) \\ &= b^{\frac{1}{y} \int_0^y \log_b f(x) \, dx} \\ &= b^{\frac{1}{y} \int_0^y \log_b b^x \, dx} \\ &= b^{\frac{1}{y} \int_0^y x \, dx} \\ &= b^{\frac{1}{y} \cdot \frac{y^2}{2}} \\ &= b^{\frac{y}{2}} \\ &= \sqrt{b^y}. \end{aligned}$$

4. Since $F(y) = \sqrt{f(y)}$, we notice that $f(y) = F(y)^2 = \exp\left(\frac{2}{y} \int_0^y \ln f(x) \, dx\right)$, and therefore $\ln f(y) = \frac{2}{y} \int_0^y \ln f(x) \, dx$.

We substitute $g(y) = \ln f(y)$, and therefore

$$yg(y) = 2 \int_0^y g(y) \, dx.$$

Therefore, differentiating both sides with respect to y gives us

$$yg'(y) + g(y) = 2g(y),$$

and therefore

$$-g(y) + yg'(y) = 0.$$

Multiplying y^{-2} on both sides gives us

$$-y^{-2}g(y) + y^{-1}g'(y) = 0,$$

and therefore

$$\frac{d}{dy} \frac{g(y)}{y} = 0,$$

and therefore

$$\frac{g(y)}{y} = C \implies g(y) = Cy.$$

Therefore, we have

$$\begin{aligned} f(y) &= \exp g(y) \\ &= \exp(Cy) \\ &= b^y \end{aligned}$$

if we substitute $b = \exp(C) > 0$, and therefore $f(x) = b^x$ as desired.

2017.3 Question 5

Since we have $x = r \cos \theta$ and $y = r \sin \theta$, and $r = f(\theta)$, we have

$$\begin{aligned}\frac{dx}{d\theta} &= \frac{dr}{d\theta} \cdot \cos \theta + r \cdot \frac{d \cos \theta}{d\theta} \\ &= f'(\theta) \cos \theta - f(\theta) \sin \theta,\end{aligned}$$

and

$$\begin{aligned}\frac{dy}{d\theta} &= \frac{dr}{d\theta} \cdot \sin \theta + r \cdot \frac{d \sin \theta}{d\theta} \\ &= f'(\theta) \sin \theta + f(\theta) \cos \theta,\end{aligned}$$

Therefore,

$$\begin{aligned}\frac{dy}{dx} &= \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} \\ &= \frac{f'(\theta) \sin \theta + f(\theta) \cos \theta}{f'(\theta) \cos \theta - f(\theta) \sin \theta} \\ &= \frac{f'(\theta) \tan \theta + f(\theta)}{f'(\theta) - f(\theta) \tan \theta}.\end{aligned}$$

For the two curves, we must have

$$\left. \frac{dy}{dx} \right|_f \cdot \left. \frac{dy}{dx} \right|_g = -1$$

for them to meet at right angles. Therefore,

$$\begin{aligned}\frac{f'(\theta) \tan \theta + f(\theta)}{f'(\theta) - f(\theta) \tan \theta} \cdot \frac{g'(\theta) \tan \theta + g(\theta)}{g'(\theta) - g(\theta) \tan \theta} &= -1 \\ (f'(\theta) \tan \theta + f(\theta)) \cdot (g'(\theta) \tan \theta + g(\theta)) &= -(f'(\theta) - f(\theta) \tan \theta) \cdot (g'(\theta) - g(\theta) \tan \theta) \\ f'(\theta)g'(\theta)(1 + \tan^2 \theta) + f(\theta)g(\theta)(1 + \tan^2 \theta) &= 0 \\ f'(\theta)g'(\theta) + f(\theta)g(\theta) &= 0.\end{aligned}$$

We have $f(-\frac{\pi}{2}) = 4$. Let

$$g_a(\theta) = a(1 + \sin \theta).$$

Therefore,

$$g'_a(\theta) = a \cos \theta,$$

and we have

$$f'(\theta)(a \cos \theta) + f(\theta)a(1 + \sin \theta) = 0,$$

and therefore

$$\frac{df(\theta)}{d\theta} \cos \theta = -f(\theta)(1 + \sin \theta).$$

By separating variables we have

$$\frac{df(\theta)}{f(\theta)} = -\frac{d\theta(1 + \sin \theta)}{\cos \theta}.$$

Notice that

$$-\frac{1 + \sin \theta}{\cos \theta} = -\frac{(1 - \sin \theta)(1 + \sin \theta)}{(1 - \sin \theta) \cos \theta} = -\frac{\cos \theta}{1 - \sin \theta} = \frac{\cos \theta}{\sin \theta - 1},$$


integrating both sides gives us

$$\ln f(\theta) = \ln |\sin \theta - 1| + C = \ln(1 - \sin \theta) + C,$$

which gives

$$f(\theta) = A(1 - \sin \theta).$$

Since $f(-\frac{\pi}{2}) = 4$, we have $2A = 4$ and $A = 2$, therefore $f(\theta) = 2(1 - \sin \theta)$.



2017.3 Question 6

1. Consider the substitution $u = \frac{1}{v}$.

When $u \rightarrow 0^+$, $v \rightarrow +\infty$.

When $u = x$, $v = \frac{1}{x}$.

We also have

$$du = -\frac{1}{v^2} dv.$$

Therefore,

$$\begin{aligned} T(x) &= \int_0^x \frac{du}{1+u^2} \\ &= \int_{+\infty}^{\frac{1}{x}} -\frac{1}{v^2} \cdot \frac{1}{1+\frac{1}{v^2}} dv \\ &= \int_{\frac{1}{x}}^{+\infty} \frac{dv}{1+v^2} \\ &= \int_0^{+\infty} \frac{dv}{1+v^2} - \int_0^{\frac{1}{x}} \frac{dv}{1+v^2} \\ &= T_\infty - T(x^{-1}), \end{aligned}$$

as desired.

2. When $u \neq a^{-1}$, we have

$$\begin{aligned} \frac{dv}{du} &= \frac{d}{du} \frac{u+a}{1-au} \\ &= \frac{1 \cdot (1-au) + a \cdot (u+a)}{(1-au)^2} \\ &= \frac{1-au+au+a^2}{(1-au)^2} \\ &= \frac{1+a^2}{(1-au)^2}. \end{aligned}$$

Also, notice that

$$\begin{aligned} \frac{1+v^2}{1+u^2} &= \frac{1 + \left(\frac{u+a}{1-au}\right)^2}{1+u^2} \\ &= \frac{(1-au)^2 + (u+a)^2}{(1+u^2)(1-au)^2} \\ &= \frac{1-2au+a^2u^2+u^2+2au+a^2}{(1+u^2)(1-au)^2} \\ &= \frac{(1+a^2)(1+u^2)}{(1-au)^2(1+u^2)} \\ &= \frac{1+a^2}{(1-au)^2}. \end{aligned}$$

Therefore, $\frac{dv}{du} = \frac{1+v^2}{1+u^2}$ as desired.

Consider the substitution $v = \frac{u+a}{1-au}$. When $u = 0$, $v = a$. When $u = x$, $v = \frac{x+a}{1-ax}$. Therefore,

$$\begin{aligned} T(x) &= \int_0^x \frac{du}{1+u^2} \\ &= \int_a^{\frac{x+a}{1-ax}} \frac{1+u^2}{1+v^2} \cdot \frac{dv}{1+u^2} \\ &= \int_a^{\frac{x+a}{1-ax}} \frac{dv}{1+v^2} \\ &= \int_0^{\frac{x+a}{1-ax}} \frac{dv}{1+v^2} - \int_0^a \frac{dv}{1+v^2} \\ &= T\left(\frac{x+a}{1-ax}\right) - T(a), \end{aligned}$$

as desired.

If we substitute $T(x) = T_\infty - T(x^{-1})$ and $T(a) = T_\infty - T(a^{-1})$, we can see that

$$\begin{aligned} T(x) &= T\left(\frac{x+a}{1-ax}\right) - T(a) \\ T_\infty - T(x^{-1}) &= T\left(\frac{x+a}{1-ax}\right) - [T_\infty - T(a^{-1})] \\ T(x^{-1}) &= 2T_\infty - T\left(\frac{x+a}{1-ax}\right) - T(a^{-1}), \end{aligned}$$

as desired.

Now, let $y = x^{-1}$ and $b = a^{-1}$. Then

$$\begin{aligned} \frac{x+a}{1-ax} &= \frac{y^{-1}+b^{-1}}{1-b^{-1}y^{-1}} \\ &= \frac{b+y}{by-1}. \end{aligned}$$

This gives us

$$T(y) = 2T_\infty - T\left(\frac{b+y}{by-1}\right) - T(b),$$

as desired.

3. Let $y = b = \sqrt{3}$. We can easily verify that $b > 0$ and $y > \frac{1}{b}$. Therefore,

$$T(\sqrt{3}) = 2T_\infty - T\left(\frac{\sqrt{3}+\sqrt{3}}{3-1}\right) - T(\sqrt{3}),$$

which simplified, gives us $T(\sqrt{3}) = \frac{2}{3}T_\infty$ as desired.

In $T(x) = T\left(\frac{x+a}{1-ax}\right) - T(a)$, let $x = a = \sqrt{2} - 1$, we can verify that $a > 0$ and $x < \frac{1}{a}$, therefore we have

$$\begin{aligned} T(\sqrt{2}-1) &= T\left(\frac{(\sqrt{2}-1)+(\sqrt{2}-1)}{1-(\sqrt{2}-1)\cdot(\sqrt{2}-1)}\right) - T(\sqrt{2}-1), \\ T(\sqrt{2}-1) &= T\left(\frac{2\sqrt{2}-2}{1-(2+1-2\sqrt{2})}\right) - T(\sqrt{2}-1), \\ T(\sqrt{2}-1) &= T\left(\frac{2\sqrt{2}-2}{2\sqrt{2}-2}\right) - T(\sqrt{2}-1), \\ 2T(\sqrt{2}-1) &= T(1). \end{aligned}$$

In $T(x) = T_\infty - T(x^{-1})$, let $x = 1$. We have

$$\begin{aligned}T(1) &= T_\infty - T(1), \\2T(1) &= T_\infty.\end{aligned}$$

Therefore, $T(\sqrt{2} - 1) = \frac{1}{4}T_\infty$, as desired.

2017.3 Question 7

$$\begin{aligned}
\frac{x^2}{a^2} + \frac{y^2}{b^2} &= \left(\frac{1-t^2}{1+t^2}\right)^2 + \left(\frac{2t}{1+t^2}\right)^2 \\
&= \frac{(1-t^2)^2 + (2t)^2}{(1+t^2)^2} \\
&= \frac{1-2t^2+t^4+4t^2}{(1+t^2)^2} \\
&= \frac{1+2t^2+t^4}{(1+t^2)^2} \\
&= \frac{(1+t^2)^2}{(1+t^2)^2} \\
&= 1
\end{aligned}$$

as desired, so T lies on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$.

1. The gradient of L must satisfy that

$$\begin{aligned}
\frac{dy}{dx} &= \frac{dy/dt}{dx/dt} \\
&= \frac{b}{a} \cdot \frac{d(2t/(1+t^2))/dt}{d((1-t^2)/(1+t^2))/dt} \\
&= \frac{b}{a} \cdot \frac{2 \cdot (1+t^2) - 2t \cdot 2t}{-2t \cdot (1+t^2) - (1-t^2) \cdot 2t} \\
&= \frac{b}{a} \cdot \frac{2+2t^2-4t^2}{-2t-2t^3-2t+2t^3} \\
&= \frac{b}{a} \cdot \frac{1-t^2}{-2t}.
\end{aligned}$$

Therefore, we have a general point $(X, Y) \in L$ satisfy that

$$\begin{aligned}
Y - \frac{2bt}{1+t^2} &= \frac{b}{a} \cdot \frac{1-t^2}{-2t} \cdot \left(X - \frac{a(1-t^2)}{1+t^2}\right) \\
(1+t^2)Y - 2bt &= \frac{b}{a} \cdot \frac{1-t^2}{-2t} \cdot ((1+t^2)X - a(1-t^2)) \\
(-2at)(1+t^2)Y - (-2at)(2bt) &= b \cdot (1-t^2) \cdot ((1+t^2)X - a(1-t^2)) \\
(-2at)(1+t^2)Y &= b(1-t^2)(1+t^2)X - ab(1-t^2)^2 - 4abt^2 \\
(-2at)(1+t^2)Y &= b(1-t^2)(1+t^2)X - ab(1+t^2)^2 \\
-2atY &= b(1-t^2)X - ab(1+t^2) \\
ab(1+t^2) - 2atY - b(1-t^2)X &= 0 \\
(a+X)bt^2 - 2aYt + b(a-X) &= 0
\end{aligned}$$

as desired.

Now if we fix X, Y and solve for t , there are two solutions to this quadratic equation exactly when

$$\begin{aligned}
(2aY)^2 - 4(a+X)b \cdot b(a-X) &> 0 \\
(aY)^2 - (a+X)(a-X)b^2 &> 0 \\
a^2Y^2 &> (a^2 - X^2)b^2,
\end{aligned}$$


which corresponds to two distinct points on the ellipse.

Since $a^2Y^2 > (a^2 - X^2)b^2$, we have $\frac{Y^2}{b^2} > 1 - \frac{X^2}{a^2}$ by dividing through a^2b^2 on both sides, i.e.

$$\frac{X^2}{a^2} + \frac{Y^2}{b^2} > 1,$$

which means when the point (X, Y) lies outside the ellipse.

This also holds when $X^2 = a^2$, i.e. when the point (X, Y) lies on the pair of lines $X = \pm A$. Here, the condition is simply $a^2Y^2 > 0$, which gives $Y \neq 0$. One of the tangents will be the vertical line $X = \pm A$ (whichever one the point lies on), and the other one as a non-vertical (as shown when $X = a$, the tangents being L_1 and L_2).



2. By Vieta's Theorem, we have

$$pq = \frac{b(a - X)}{b(a + X)} \implies (a + X)pq = a - X,$$

as desired, and

$$p + q = -\frac{-2aY}{(a + X)b} = \frac{2aY}{(a + X)b}.$$

Let $X = 0$ for the equation in L ,

$$abt^2 - 2aYt + ba = 0$$

$$bt^2 - 2Yt + b = 0$$

$$Y = \frac{b(1 + t^2)}{2t}.$$

Therefore,

$$\begin{aligned} y_1 + y_2 &= \frac{b(1 + p^2)}{2p} + \frac{b(1 + q^2)}{2q} \\ &= \frac{b[(1 + p^2)q + (1 + q^2)p]}{2pq} \\ &= 2b, \end{aligned}$$

therefore we have

$$4pq = (1 + p^2)q + (1 + q^2)p = (p + q)(1 + pq).$$

Therefore,

$$\begin{aligned}4 \cdot \frac{a-X}{a+X} &= \frac{2aY}{(a+X)b} \cdot \frac{2a}{a+X} \\ a-X &= \frac{a^2Y}{b(a+X)} \\ (a-X)(a+X)b &= a^2Y \\ (a^2-X^2)b &= a^2Y \\ 1 - \frac{X^2}{a^2} &= \frac{Y}{b} \\ \frac{X^2}{a^2} + \frac{Y}{b} &= 1,\end{aligned}$$

as desired.

2017.3 Question 8

We have

$$\begin{aligned}
 \sum_{m=1}^n a_m(b_{m+1} - b_m) &= \sum_{m=1}^n a_m b_{m+1} - \sum_{m=1}^n a_m b_m \\
 &= -\sum_{m=0}^{n-1} b_{m+1} a_{m+1} + \sum_{m=1}^n b_{m+1} a_m \\
 &= -\sum_{m=1}^n b_{m+1} a_{m+1} + \sum_{m=1}^n b_{m+1} a_m + a_{n+1} b_{n+1} - a_1 b_1 \\
 &= a_{n+1} b_{n+1} - a_1 b_1 - \sum_{m=1}^n b_{m+1} (a_{m+1} - a_m),
 \end{aligned}$$

as desired.

1. Let $a_m = 1$. On one hand, we have

$$\begin{aligned}
 \sum_{m=1}^n a_m(b_{m+1} - b_m) &= \sum_{m=1}^n [\sin(m+1)x - \sin mx] \\
 &= \sum_{m=1}^n 2 \cos\left(\frac{(m+1)x + mx}{2}\right) \sin\left(\frac{(m+1)x - mx}{2}\right) \\
 &= 2 \sum_{m=1}^n \cos\left(m + \frac{1}{2}\right)x \sin \frac{x}{2} \\
 &= 2 \sin \frac{x}{2} \sum_{m=1}^n \cos\left(m + \frac{1}{2}\right)x.
 \end{aligned}$$

On the other hand, we have

$$\begin{aligned}
 \sum_{m=1}^n a_m(b_{m+1} - b_m) &= a_{n+1} b_{n+1} - a_1 b_1 - \sum_{m=1}^n b_{m+1} (a_{m+1} - a_m) \\
 &= \sin(n+1)x - \sin x.
 \end{aligned}$$

Therefore, by rearranging, we have

$$\sum_{m=1}^n \cos\left(m + \frac{1}{2}\right)x = \frac{1}{2} [\sin(n+1)x - \sin x] \operatorname{cosec} \frac{1}{2}x$$

as desired.

2. Let $a_m = m$, and let $b_m = \cos\left(m - \frac{1}{2}\right)x$. We have the identity

$$\cos A - \cos B = -2 \sin\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right).$$

Therefore, we have

$$\begin{aligned}
 \sum_{m=1}^n a_m(b_{m+1} - b_m) &= \sum_{m=1}^n m \cdot \left[\cos\left(m + \frac{1}{2}\right)x - \cos\left(m - \frac{1}{2}\right)x \right] \\
 &= \sum_{m=1}^n -2m \sin mx \sin \frac{1}{2}x \\
 &= -2 \sin \frac{1}{2}x \sum_{m=1}^n m \sin mx,
 \end{aligned}$$

and

$$\begin{aligned}
& \sum_{m=1}^n a_m(b_{m+1} - b_m) \\
&= a_{n+1}b_{n+1} - a_1b_1 - \sum_{m=1}^n b_{m+1}(a_{m+1} - a_m) \\
&= (n+1) \cos\left(n + \frac{1}{2}\right)x - 1 \cdot \cos\frac{1}{2}x - \sum_{m=1}^n \cos\left(m + \frac{1}{2}\right)x \cdot 1 \\
&= (n+1) \cos\left(n + \frac{1}{2}\right)x - \cos\frac{1}{2}x - \sum_{m=1}^n \cos\left(m + \frac{1}{2}\right)x \\
&= (n+1) \cos\left(n + \frac{1}{2}\right)x - \cos\frac{1}{2}x - \frac{1}{2}(\sin(n+1)x - \sin x) \operatorname{cosec}\frac{1}{2}x \\
&= \frac{1}{2} \operatorname{cosec}\frac{1}{2}x \left[2(n+1) \cos\left(n + \frac{1}{2}\right)x \sin\frac{1}{2}x - 2 \cos\frac{1}{2}x \sin\frac{1}{2}x - (\sin(n+1)x - \sin x) \right] \\
&= \frac{1}{2} \operatorname{cosec}\frac{1}{2}x [(n+1)(\sin(n+1)x - \sin nx) - (\sin x - \sin 0) - (\sin(n+1)x - \sin x)] \\
&= \frac{1}{2} \operatorname{cosec}\frac{1}{2}x [n \sin(n+1)x - (n+1) \sin nx].
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
-2 \sin\frac{1}{2}x \sum_{m=1}^n m \sin mx &= \frac{1}{2} \operatorname{cosec}\frac{1}{2}x [n \sin(n+1)x - (n+1) \sin nx] \\
\sum_{m=1}^n m \sin mx &= -\frac{1}{4} \operatorname{cosec}^2\frac{1}{2}x [n \sin(n+1)x - (n+1) \sin nx],
\end{aligned}$$

and therefore, $p = -\frac{1}{4}n$, $q = \frac{1}{4}(n+1)$.

2017.3 Question 12

1. First, note that

$$\begin{aligned}
 1 &= \sum_{x,y=1}^{x=n} P(X = x, Y = y) \\
 &= \sum_{x=1}^n \sum_{y=1}^n k(x + y) \\
 &= \sum_{x=1}^n \sum_{y=1}^n (kx + ky) \\
 &= \sum_{x=1}^n \left(n \cdot kx + k \sum_{y=1}^n y \right) \\
 &= nk \sum_{x=1}^n x + nk \sum_{y=1}^n y \\
 &= n^2(n + 1)k
 \end{aligned}$$

Therefore, $k = \frac{1}{n^2(n+1)}$

$$\begin{aligned}
 P(X = x) &= \sum_{y=1}^n P(X = x, Y = y) \\
 &= \sum_{y=1}^n k(x + y) \\
 &= nkx + k \sum_{y=1}^n y \\
 &= nkx + \frac{kn(n + 1)}{2} \\
 &= \frac{x}{n(n + 1)} + \frac{1}{2n} \\
 &= \frac{2x + n + 1}{2n(n + 1)},
 \end{aligned}$$

as desired.

By symmetry, $P(Y = y) = \frac{2y+n+1}{2n(n+1)}$.

We have

$$P(X = x) \cdot P(Y = y) = \frac{(2x + n + 1)(2y + n + 1)}{4n^2(n + 1)^2}.$$

But $P(X = x, Y = y) = \frac{x+y}{n^2(n+1)}$ is not equal to this. So X and Y are not independent.

2. By definition,

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y).$$

We have

$$\begin{aligned}
 E(X) = E(Y) &= \sum_{t=1}^n t \cdot P(X = t) \\
 &= \sum_{t=1}^n \frac{t \cdot (2t + n + 1)}{2n(n + 1)} \\
 &= \frac{1}{n(n + 1)} \sum_{t=1}^n t^2 + \frac{1}{2n} \sum_{t=1}^n t \\
 &= \frac{n(n + 1)(2n + 1)}{6n(n + 1)} + \frac{n(n + 1)}{4n} \\
 &= \frac{2n + 1}{6} + \frac{n + 1}{4} \\
 &= \frac{4n + 2 + 3n + 3}{12} \\
 &= \frac{7n + 5}{12},
 \end{aligned}$$

and

$$\begin{aligned}
 E(XY) &= \sum_{x,y=1}^n xy \cdot P(X = x, Y = y) \\
 &= \sum_{x=1}^n \sum_{y=1}^n \frac{xy(x + y)}{n^2(n + 1)} \\
 &= \frac{1}{n^2(n + 1)} \sum_{x=1}^n \sum_{y=1}^n xy(x + y) \\
 &= \frac{1}{n^2(n + 1)} \sum_{x=1}^n \sum_{y=1}^n (x^2y + xy^2) \\
 &= \frac{1}{n^2(n + 1)} \left[\sum_{x=1}^n x^2 \sum_{y=1}^n y + \sum_{x=1}^n x \sum_{y=1}^n y^2 \right] \\
 &= \frac{1}{n^2(n + 1)} \cdot 2 \cdot \frac{n(n + 1)(2n + 1)}{6} \cdot \frac{n(n + 1)}{2} \\
 &= \frac{(2n + 1)(n + 1)}{6}.
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 \text{Cov}(X, Y) &= E(XY) - E(X)E(Y) \\
 &= \frac{(2n + 1)(n + 1)}{6} - \frac{(7n + 5)^2}{144} \\
 &= \frac{48n^2 + 72n + 24}{144} - \frac{49n^2 + 70n + 25}{144} \\
 &= \frac{-n^2 + 2n - 1}{144} \\
 &= -\frac{(n - 1)^2}{144} \\
 &< 0,
 \end{aligned}$$

as desired.

2017.3 Question 13

We have

$$\begin{aligned} V(x) &= E[(X - x)^2] \\ &= E(X^2 - 2xX + x^2) \\ &= E(X^2) - 2xE(X) + x^2 \\ &= \sigma^2 + \mu^2 - 2x\mu + x^2. \end{aligned}$$

Therefore, if $Y = V(X)$, then

$$\begin{aligned} E(Y) &= E(V(X)) \\ &= E(\sigma^2 + \mu^2 - 2X\mu + X^2) \\ &= \sigma^2 + \mu^2 - 2\mu E(X) + E(X^2) \\ &= \sigma^2 + \mu^2 - 2\mu^2 + \mu^2 + \sigma^2 \\ &= 2\sigma^2. \end{aligned}$$

Let $X \sim U[0, 1]$, we have $\mu = E(X) = \frac{1}{2}$, and $\sigma^2 = \text{Var}(X) = \frac{1}{12}$. Therefore,

$$\begin{aligned} V(x) &= \frac{1}{12} + \frac{1}{4} - x + x^2 \\ &= x^2 - x + \frac{1}{3}. \end{aligned}$$

The c.d.f. of X is F , defined as

$$P(X \leq x) = F(x) = \begin{cases} 0, & x \leq 0, \\ x, & 0 < x \leq 1, \\ 1, & 1 < x \end{cases}$$

Let the c.d.f. of Y be G , we have $G(y) = P(Y \leq y)$.

Since $V([0, 1]) = [\frac{1}{12}, \frac{1}{3}]$, we must have $G(y) = 0$ for $y \leq \frac{1}{12}$ and $G(y) = 1$ for $y > \frac{1}{3}$.

For $y \in (\frac{1}{12}, \frac{1}{3}]$, we have

$$\begin{aligned} G(y) &= P(Y \leq y) = P(V(X) \leq y) \\ &= P\left(\left(x - \frac{1}{2}\right)^2 + \frac{1}{12} \leq y\right) \\ &= P\left(\left|x - \frac{1}{2}\right| \leq \sqrt{y - \frac{1}{12}}\right) \\ &= P\left(\frac{1}{2} - \sqrt{y - \frac{1}{12}} \leq x \leq \frac{1}{2} + \sqrt{y - \frac{1}{12}}\right) \\ &= F\left(\frac{1}{2} + \sqrt{y - \frac{1}{12}}\right) - F\left(\frac{1}{2} - \sqrt{y - \frac{1}{12}}\right) \\ &= \left(\frac{1}{2} + \sqrt{y - \frac{1}{12}}\right) - \left(\frac{1}{2} - \sqrt{y - \frac{1}{12}}\right) \\ &= 2\sqrt{y - \frac{1}{12}}. \end{aligned}$$

Therefore, the p.d.f. of y , g satisfies that for $y \in (\frac{1}{12}, \frac{1}{3}]$,

$$g(y) = G'(y) = \frac{1}{\sqrt{y - \frac{1}{12}}}$$

and 0 everywhere else.